

Boston University College of Health
& Rehabilitation Sciences: Sargent College
Center for Psychiatric Rehabilitation



940 Commonwealth Avenue, West
Boston, Massachusetts 02215
T 617-353-3549 F 617-353-7700
www.bu.edu/cpr

January 14, 2026

The Honorable Michael Day
Chair, Joint Committee on Judiciary
24 Beacon Street, Room 136
Boston, MA 02133
Michael.day@mahouse.gov
Talia.Quinn@mahouse.gov

Re: Testimony in opposition to H. 1801, *An Act to provide continuum of care for severe mental illness*

Dear Chair Day and House Members of the Committee on the Judiciary:

My name is Kim Mueser and I am a clinical psychologist and professor in the Departments of Occupational Therapy and Psychological and Brain Sciences at Boston University. I was formally the Executive Director of the Center for Psychiatric Rehabilitation at Boston University, where I still work. My area of research is on the development and evaluation of psychosocial interventions for people with schizophrenia and other serious mental illnesses (SMI), and more broadly on evidence-based practices for the treatment of SMI. I am the co-author of over 400 peer reviewed journal articles, over 20 books, and over 100 book chapters. I have also had family members with SMI.

An “evidence-based practice” is an intervention for a disorder that has been scientifically evaluated in multiple rigorously controlled studies and found to improve outcomes more than standard treatment.¹ I oppose the passage of this bill allowing “assisted outpatient treatment” (AOT) or commitment to outpatient mental health treatment of people with SMI for the simple reason that research indicates that it does not work, and the broad consensus in the field at this time is that AOT is *not* an evidence-based practice for improving the outcome of SMI.^{2,3}

While AOT is not effective, the good news is that a growing number of evidence-based practices have been shown to be effective, including several psychosocial interventions. I will briefly describe four of these interventions here.

First, training in *illness management and recovery* involves teaching individuals with SMI about the nature of their disorder and how to manage it while helping them set and pursue their own recovery goals. Teaching clients such information and skills empowers them to take control over their illness and to get on with their lives. Indeed, research shows that illness

management and recovery programs reduce hospitalizations and improve psychosocial functioning and quality of life in people with SMI.⁴⁻⁶

Second, *family psychoeducation* is an intervention in which a member of the client's treatment team works with the family (including the client) to teach them about psychiatric illness and the principles of its management, as well as strategies for reducing stress and solving problems together. By reaching out and engaging family members in a collaborative relationship with the treatment team, family psychoeducation is effective at reducing relapses and rehospitalizations, lowering family burden, and increasing client functioning.⁷⁻⁹

Third, *supported employment* is an approach to helping individuals with SMI get and keep competitive jobs in areas that are of interest to them. In contrast to traditional vocational rehabilitation programs in which individuals are given extensive amounts of training before helping them find jobs, supported employment works by rapid job search for competitive jobs (usually within a month of joining a program), and then providing supports to help the person stay on and succeed in the job. Numerous studies show that supported employment is both more effective at improving competitive work outcomes than other approaches,^{10,11} and that success in achieving work is associated with significant reductions in psychiatric hospitalizations.¹²⁻¹⁴

Fourth and last, Coordinated Specialty Care programs (CSC) are for people who have recently developed a first episode of psychosis, which is often the beginning of a schizophrenia spectrum disorder.¹⁵ CSC programs typically provide pharmacological treatment, family, psychoeducation, individual or group psychotherapy, supported employment and education, and peer support. Extensive research throughout the world shows that CSC programs improve the broad range of outcomes of individuals early in the treatment of a psychosis, including symptom severity, psychosocial functioning, involvement in work or school, while also reducing psychiatric hospitalizations.¹⁶

We have numerous tools at our disposal in the form of evidence-based practices for improving the long-term course of SMI. We owe it to individuals with SMI to provide them with the most effective treatments possible based on the best available scientific research, and to refrain from imposing interventions that lack empirical support, especially those that limit individual freedoms and are potentially harmful, such as outpatient commitment to treatment.


References

1. Drake RE, Goldman HH, Leff HS, et al. Implementing evidence-based practices in routine mental health service settings. *Psychiatric Services*. 2001;52:179-182.
2. Kisely SR, Campbell LA, O'Reilly R. Compulsory community and involuntary outpatient treatment for people with severe mental disorders. *Cochrane Database of Systematic Reviews*. 2017;3:CD004408. doi:10.1002/14651858.CD004408.pub5
3. Segal SP. Protecting health and safety with needed-treatment: The effectiveness of outpatient commitment. *Psychiatric Quarterly*. 2022;93:55-79.
4. Barnicot K, Michael C, Trione E, et al. Psychological interventions for acute

psychiatric inpatients with schizophrenia-spectrum disorders: A systematic review and meta-analysis. *Clinical Psychology Review*. 2020;82:101929

5. Bighelli I, Rodolico A, García-Mieres H, et al. Psychosocial and psychological interventions for relapse prevention in schizophrenia: a systematic review and network meta-analysis. *Lancet Psychiatry*. 2021;8:969-980. doi:10.1016/S2215-0366(21)00243-1
6. Lincoln TM, Wilhelma K, Nestoriuca Y. Effectiveness of psychoeducation for relapse, symptoms, knowledge, adherence and functioning in psychotic disorders: A meta-analysis. *Schizophrenia Research*. 2007;96:232-245.
7. Camacho-Gomez M, Castellvi P. Effectiveness of family intervention for preventing relapse in first-episode psychosis until 24 months of follow-up: A systematic review with meta-analysis of randomized controlled trials. *Schizophrenia Bulletin*. 2020;46:98-109. doi:10.1093/schbul/sbz038
8. Rodolico A, Bighelli I, Avanzato C, et al. Family interventions for relapse prevention in schizophrenia: A systematic review and network meta-analysis. *Lancet Psychiatry*. 2022;9:211-221. doi:10.1016/S2215-0366(21)00437-5
9. Yesufu-Udechuku A, Harrison B, Mayo-Wilson E, et al. Interventions to improve the experience of caring for people with severe mental illness: Systematic review and meta-analysis. *British Journal of Psychiatry*. 2015;206:268-74. doi:10.1192/bjp.bp.114.147561
10. Bond GR, Al-Abdulmunem M, Marbacher J, Christensen TN, Sveinsdottir V, Drake RE. A systematic review and meta-analysis of IPS supported employment for young adults with mental health conditions. *Administration & Policy in Mental Health*. 2023;160–172:160–172. doi:10.1007/s10488-022-01228-9
11. Frederick DE, VanderWeele TJ. Supported employment: Meta-analysis and review of randomized controlled trials of individual placement and support. *PLoS ONE*. 2019;14(2):e0212208.
12. Burns T, Caty J, White S, et al. The impact of supported employment and working on clinical and social functioning: Results of an international study of individual placement and support. *Schizophrenia Bulletin*. 2009;35:949-958.
13. Hoffmann H, Jäckel D, Glauser S, Mueser KT, Kupper Z. Long-term effectiveness of supported employment: Five-year follow-up of a randomized controlled trial. *American Journal of Psychiatry*. 2014;171:1183-1190. doi:10.1176/appi.ajp.2014.13070857
14. Mueser KT, Becker DR, Torrey WC, et al. Work and nonvocational domains of functioning in persons with severe mental illness: A longitudinal analysis. *Journal of Nervous and Mental Disease*. 1997;185:419-426.
15. Heinsen RK, Goldstein AB, Azrin ST. Evidence-Based Treatments for First Episode Psychosis: Components of Coordinated Specialty Care. National Institute of Mental Health; 2014.
16. Correll CU, Galling B, Pawar A, et al. Comparison of early intervention services vs treatment as usual for early-phase psychosis: A systematic review, meta-analysis, and meta-regression. *JAMA Psychiatry*. 2018;75:555-65. doi:10.1001/jamapsychiatry.2018.0623

Sincerely,

A handwritten signature in black ink, appearing to read "Kim T. Mueser". The signature is fluid and cursive, with a long horizontal stroke at the end.

Kim T. Mueser, Ph.D.
Center for Psychiatric Rehabilitation
Professor
Departments of Occupational Therapy and Psychological & Brain Sciences
Boston University

Phone: 617-353-3549

FAX: 617-353-7700

Email: mueser@bu.edu